

A Serendipitous Search for High-Redshift Ly α Emission: Two Primeval Galaxy Candidates at $z \simeq 3$ ¹

Curtis Manning, Daniel Stern², Hyron Spinrad, & Andrew J. Bunker³

Department of Astronomy, University of California at Berkeley

Berkeley, CA 94720

email: (cmanning,dstern,spinrad,bunker)@bigz.berkeley.edu

ABSTRACT

In the course of our ongoing search for serendipitous high-redshift Ly α emission in deep archival Keck spectra, we discovered two very high equivalent width ($W_{\lambda}^{\text{obs}} \gtrsim 450$ Å, 2σ) Ly α emission line candidates at $z \sim 3$ in a moderate dispersion ($\lambda/\Delta\lambda \simeq 1200$) spectrogram. Both lines have low velocity dispersions ($\sigma_v \sim 60$ km s⁻¹) and deconvolved radii $r \approx 1 h_{50}^{-1}$ kpc. We argue that the lines are Ly α , and are powered by stellar ionization. The surface density of robust, high equivalent width Ly α candidates is estimated to be $\sim 3 \pm 2$ arcmin⁻² per unit redshift at $z \simeq 3$, consistent with the estimate of Cowie & Hu (1998). The Ly α emission line source characteristics are consistent with the galaxies undergoing their first burst of star formation, i.e., with being primeval. Source sizes and velocity dispersions are comparable to the theoretical primeval galaxy model of Lin & Murray (1992) based on the inside-out, self-similar collapse of an isothermal sphere. In this model, star formation among field galaxies is a protracted process.

Galaxies are thought to be able to display high equivalent widths for only the first $\sim \text{few} \times 10^7$ yr. This time is short in relation to the difference in look back times between $z = 3$ and $z = 4$, and implies that a substantial fraction of strong line-emitting galaxies at $z = 3$ were formed at redshifts $z \leq 4$. We discuss the significance of high-equivalent width Ly α -emitting galaxies in terms of the emerging picture of the environment, and the specific characteristics of primeval galaxy formation at high redshift.

Subject headings: Cosmology: observations — galaxies: compact — galaxies: evolution — galaxies:formation — galaxies: starburst — line: profiles

¹Based on observations at the W.M. Keck Observatory, which is operated as a scientific partnership among the University of California, the California Institute of Technology, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

²Current address: Jet Propulsion Laboratory, California Institute of Technology, Mail Stop 169-327, Pasadena, CA 91109; stern@zwoolfkinder.jpl.nasa.gov

³Current address: Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, England; email: bunker@ast.cam.ac.uk

1. Introduction

Early theoretical models of primeval galaxies suggested that their Ly α lines would be highly luminous and diffuse. But searches on 4m-class telescopes confirmed no such emission lines (Pritchet 1994; Thompson & Djorgovski 1995) and the search for galaxies at large look-back times turned to other strategies. One highly successful technique is color-selection of the Lyman break in galaxies of redshift $z \simeq 3$ (Steidel & Hamilton 1992; Steidel et al. 1996). Lyman-break galaxies (LBGs) are strongly star-forming, and compact relative to local $\mathcal{L}^*$ galaxies (Giavalisco et al. 1996b). However, LBGs are not generally thought to be primeval, in part because they have lower Ly α equivalent widths than expected of primeval galaxies, in part because interstellar metallic absorption lines (e.g., Si II λ 1260, Al II λ 1670) are characteristic of most LBG spectra.

The theoretical upper limit equivalent width for Ly α emission produced by stellar photoionization is thought to be $W_{\lambda}^{\text{rest}} \sim 100 - 200 \text{ \AA}$ (Charlot & Fall 1993), and can be expected to last only a few $\times 10^7$ yr for a constant star formation rate (SFR). Several examples of high-redshift, high equivalent width (non-AGN) Ly α emission have been reported (e.g., Dey et al. 1998; Weymann et al. 1998; Hu et al. 1999). Few secure examples of isolated, high equivalent widths that are not Ly α exist in the literature (e.g., Stockton & Ridgway 1998; Stern et al. 2000). About half of LBGs have Ly α in emission (Steidel et al. 1996, 1998; Lowenthal et al. 1997), with rest equivalent widths $\sim 5 - 35 \text{ \AA}$. Likewise, about half of local actively star-forming galaxies have Ly α in emission, with rest equivalent widths in the range $W_{\lambda} \simeq 10 - 35 \text{ \AA}$ (Giavalisco et al. 1996a), though a few have $W_{\lambda} \gtrsim 50 \text{ \AA}$.

Local studies show that whether Ly α is seen in emission or absorption may depend heavily on the kinematics of the surrounding neutral H I halo (Legrand et al. 1997; Kunth et al. 1998a) and the chance geometry of neutral gas and dust (Giavalisco et al. 1996a). However, $W_{\lambda}^{\text{rest}}$ is shown to be weakly-correlated with metallicity (Terlevich et al. 1993; Giavalisco et al. 1996a). The selective absorption of resonantly scattered Ly α photons, whose path length in the cloud is much longer than that of photons of modestly different wavelength, is thought to be the major factor in quenching the Ly α line. The partially extinguished Ly α line is characteristically asymmetric (e.g., Dey et al. 1998; Kunth et al. 1998a; Legrand et al. 1997), presumably caused by the absorption of photons from the blue side of the line by dust in an expanding shell, and compounded by the addition of redshifted backscattering off the neutral shell expanding away from the observer (e.g., Legrand et al. 1997). Though dust-free galaxies are the most likely to have high equivalent width Ly α emission, it is clear that a more evolved galaxy can also have high equivalent width Ly α emission emerging within specific zones of active star formation due to an incomplete covering factor. Such emission may appear spatially offset from the region of star formation (e.g., Bunker et al. 2000). Averaged over the whole galaxy, the continuum supplied by recent star formation with locally quenched Ly α emission will reduce the integrated line equivalent width.

Deep spectra taken at the Keck telescopes regularly reveal serendipitous high-equivalent width, isolated emission lines (e.g., Stern et al. 2000). Indeed, the first confirmed galaxy at $z > 5$ was

discovered in this manner (Dey et al. 1998). Steidel et al. (1998) obtained a serendipitous spectrum of a strong Ly α line with extremely faint continuum near the $\langle z \rangle \simeq 3.09$ structure in the SSA22 field. Strong emission lines can also be effectively targeted by using narrow band imaging, in conjunction with broad band imaging. Recent observations, using narrow band imaging with a “strong equivalent width” ($W_{\lambda}^{\text{obs}} \geq 77 \text{ \AA}$) criterion (Cowie & Hu 1998) and follow-up spectroscopy (Hu et al. 1998) have disclosed a population of galaxies with what is thought to be Ly α in emission. The galaxies tend to have very weak, sometimes undetectable, continua. They are also quite compact, as noted of the $z \sim 2.4$ Ly α emitters found in HST searches (Pascarelle et al. 1996; Pascarelle et al. 1998). Cowie & Hu (1998) estimate a surface density of 3.6 arcmin^{-2} per unit redshift for flux densities $j > 2 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$, corresponding to line luminosities $\mathcal{L}_{\text{Ly}\alpha} > 1.8 \times 10^{42} \text{ ergs s}^{-1}$ for an Einstein-de Sitter Universe with $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ at $z \approx 3.4$.

We are conducting a search for serendipitous emission lines in deep archival Keck spectra. Such a serendipitous search is an emission-line flux-limited survey, and may find emission lines over a wide range of redshifts. Narrow-band surveys are sensitive to candidates over only small redshift ranges, and require follow-up spectroscopy to discriminate stellar Ly α lines from metal and AGN emission lines. The identification of a population of primeval galaxies may provide important information about the epoch(s) of galaxy formation, including data relevant to the integrated global star formation rate, and the luminosity function of galaxies to photometric limits fainter than that accessible to color-selection surveys.

One drawback faced by all Ly α emission line searches is possible confusion with other isolated, high equivalent width lines such as [O II] $\lambda 3727$ and H α . Experience has shown that [O II] $\lambda 3727$ is often mis-identified as Ly α . When continuum is detected, we may use the Balmer series or the continuum depression to discriminate, and when absent we may search for another line, such as N V $\lambda 1240$ or [O III] $\lambda 5007$. However, N V $\lambda 1240$ is rarely detected in non-AGN, and the search for [O III] $\lambda 5007$ often fails because these lines would either fall out of the range of the spectrograph, or in the high-noise infrared part of the spectrum which, combined with the large intrinsic dispersion in [O II] $\lambda 3727$ /[O III] $\lambda 5007$ (Kennicutt 1992), can make an unambiguous verification untenable. Alternatively, one may use the known tendency of Ly α lines to display a P-Cygni profile to identify this resonance line. However, asymmetry cannot be a completely necessary criterion for Ly α emission lines since some local sources apparently have symmetric, or “pure” profiles (Kunth et al. 1998b). When emission lines are unresolved, and have undetected continua, the last recourse is the knowledge that [O II] $\lambda 3727$ rarely has an equivalent width greater than $100 \text{ \AA}$ (but see Stern et al. 2000). For these reasons, unambiguous Ly α lines are rare, and worthy of extra scrutiny.

We serendipitously found two isolated emission lines on a single slitlet from a mask centered on the SSA22 field, and determined that they are in fact Ly α lines — examples of galaxies which are more compact and have higher Ly α equivalent widths than known LBGs. We report on their equivalent widths, intrinsic radii, velocity dispersions, and surface/volume density in order to place them in the picture of the evolving Universe emerging from studies of high-redshift galaxies. The source targeted by the slitlet in which we found our emission line galaxies is the

color-selected Lyman-break galaxy SSA22 C17; we analyze it in parallel for comparison. We adopt $H_0 = 50 h_{50} \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_0 = 1$, and $\Lambda = 0$ unless otherwise stated. At $z = 3$, for $\Omega = 1(0.1)$, $1''.0$ corresponds to $7.3 (12.3) h_{50}^{-1} \text{ kpc}$.

2. Observations and Data Reduction

We have obtained deep, moderate-dispersion spectra of $z \simeq 3$ LBGs with the aim of making detailed studies of the ages, kinematics, dust-content, and abundances of the LBG population (Dey et. al., in preparation). The data were taken during the years 1997 to 1999 with the Low Resolution Imaging Spectrometer (Oke et al. 1995) at the Cassegrain focus on the Keck II telescope. The camera uses a Tek 2048² CCD detector with a pixel scale $0''.212 \text{ pix}^{-1}$. On UT 1997 September 10 we obtained moderate-dispersion multislit spectra of LBGs in the SSA22 field, using the 600 lines mm^{-1} grating blazed at $5000 \text{ \AA}$. Slitlet widths were $1''.25$, resulting in a spectral resolution of $\sim 4.4 \text{ \AA}$ (FWHM) for sources filling the slitlet, and a resolution $\lambda/\Delta\lambda_{\text{FWHM}} \sim 1200$. Objects not filling the slitlet will have higher spectral resolution. For a spatially unresolved source – one blurred only by atmospheric seeing – we estimate a resolution of $\sim 2.8 \text{ \AA}$ (FWHM) based on Ly α absorption systems in the quasar SSA22 D14 also observed on the slitmask reported herein. Four of the integrations totaling 7200s were of excellent quality, with seeing along the spatial axis of $0''.78$. Additional observations were made on UT 1997 September 12 and UT 1999 June 14. Here we concentrate on the data of UT 1997 September 10 for which the signal-to-noise ratio (S/N) is highest.

The data reductions were performed using the IRAF package, and followed standard slit spectroscopy procedures. Flat-fielding, sky subtraction, cosmic ray removal, and aperture extractions of the slitmask data were facilitated by the home-grown software package BOGUS, created by D. Stern, A. J. Bunker & S. A. Stanford. Wavelength calibration was performed using a HgNeAr lamp, employing telluric sky lines to adjust the wavelength zero-point to the data frames, compensating for any drift in the wavelength coverage. Flux calibration was performed using observations of Wolf 1346 (Massey & Gronwall 1990).

3. Spectroscopic Results

We find two serendipitous, isolated emission lines, ser-1 and ser-2, in a $100''$ long slitlet centered on the LBG SSA22 C17 ($z = 3.299$; see Fig. 1). In lower dispersion spectra, it is often impossible to distinguish between high equivalent width forms of [O II] $\lambda 3727$ and Ly α , unless there is an evident continuum depression, a line asymmetry, or other distinguishing features. Though only one of these two serendipitously discovered sources (ser-2) displays evidence of the asymmetry characteristic of high-redshift Ly α emission in LBGs, we argue that Ly α is indeed the most likely interpretation for both. We simulate the [O II] $\lambda\lambda 3726, 3729$ doublet using our $2.8 \text{ \AA}$ resolution




Fig. 1.— Two-dimensional spectrogram of the slitlet in the SSA22 field targeting the LBG SSA22 C17 at $z = 3.299$. Two strong line emitters are serendipitously identified, ser-1 and ser-2. The high-equivalent widths, narrow velocity widths, and lack of secondary spectral features strongly argue that these are Ly α emitters at $z \simeq 3$. Note the foreground continuum source and the residual of the [O I] λ 5577 Å skyline.

(for an unresolved source) with 50 km s^{-1} intrinsic velocity dispersion, introducing the Poisson noise appropriate to the observed flux of the median of ser-1 and ser-2 (see Fig. 2). We conclude that [O II] $\lambda\lambda 3726, 3729$ is inconsistent with our lines, as the oxygen doublet would be marginally resolved and has a significantly greater FWHM. Further, the absence of associated emission lines argues against H β and [O III] $\lambda\lambda 4959, 5007$ interpretations. Finally, the lines are shortward of 6563 Å so H α is not a viable identification. In the following, we assume what is most certainly the case — that these are in fact Ly α emission lines. Notably, these spectra display no detectable continua.

The spectra of the serendipitous emission lines ser-1 and ser-2 are presented in Fig. 3, together with that of the LBG SSA22 C17. The redshifts of these galaxies, their emission line wavelengths, fluxes and attributed luminosities are given in Table 1. It should be noted that the fluxes and luminosities represent only that part of the atmospherically smeared image of the galaxy falling within the slitlet, and are in all likelihood under-estimates as these serendipitously-identified sources

Table 1: Emission Line Properties

Object	λ^{obs} (Å)	z	j_{-17}^a (cgs)	$\mathcal{L}_{42}^b$ (cgs)	W_{λ}^{obs} (Å)	$W_{\lambda}^{\text{rest}}$ (Å)	r^c (h_{50}^{-1} kpc)	σ_v^d (km s^{-1})
ser-1	4889.3	3.022	1.85	1.29	$\geq 550 (2\sigma)$	$\geq 137 (2\sigma)$	0.8 ± 0.6	62 ± 9
ser-2	5296.8	3.357	1.35	1.19	$\geq 470 (2\sigma)$	$\geq 109 (2\sigma)$	1.1 ± 0.5	47 ± 9
C17	5226.4	3.299	2.60	2.20	35.6	8.3	2.8:	82:

Notes.— (a) Emission line fluxes are in units of $10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$. (b) The Ly α line luminosities are in units of $10^{42} h_{50}^{-2} \text{ ergs s}^{-1}$. (c) radius based on deconvolution of source FWHM. (d) Deconvolved velocity dispersion assumes sources fill the $1''.25$ width of slitlets.




Fig. 2.— The [O II] $\lambda\lambda 3726, 3729$ doublet line profile from a source with a 50 km s^{-1} velocity dispersion, as it would appear in its rest frame, observed with an effective $2.8 \text{ \AA}$ resolution of a point-source emitter. Poisson noise has been added commensurate to that observed in the LRIS spectra. The observed emission lines of ser-1 and ser-2 have been referred to the same wavelength, with their amplitude adjusted to agree with that of the [O II] $\lambda\lambda 3726, 3729$ lines. The line profiles of ser-1 and ser-2 are clearly incompatible with an [O II] $\lambda\lambda 3726, 3729$ interpretation.

are not necessarily centered in the slitlets. The sources ser-1 and ser-2 are at redshifts $z = 3.022$, and $z = 3.357$, respectively. We note that ser-2 is at nearly the same redshift as the nearby quasar SSA22 D14 (Steidel et al. 1998).

To date, our serendipitous search has reviewed five slitmasks ($\simeq 0.75 \text{ arcmin}^2$) and five longslit spectra ($\simeq 0.30 \text{ arcmin}^2$). In addition to the two lines presented here, we have found one other definite Ly α line at $z = 2.946$ with a line flux $j = 3.32 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and line luminosity $\mathcal{L}_{\text{Ly}\alpha} = 2.2 \times 10^{42} h_{50}^{-2} \text{ ergs s}^{-1}$. In a total search area slightly over 1 arcmin^2 , this suggests that the surface density of robustly identified high equivalent width Ly α emission lines within the redshift range, $2.5 \leq z \leq 3.5$ is very roughly of the order $3 \pm 2 \text{ arcmin}^{-2}$ per unit redshift. This is consistent with the estimate of Cowie & Hu (1998) who find a surface density of $\approx 3.6 \text{ Ly}\alpha\text{-emitters arcmin}^{-2}$ per unit redshift to a higher limiting flux density of $2 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$. The full survey including candidates at higher redshift will be discussed in a subsequent paper.




Fig. 3.— Spectra of two serendipitously identified Ly α -emitters in the field of SSA22, with the known LBG SSA22 C17. The abscissa on the left refers to the first two plots, while the abscissa on the right refers to the right-hand plot. Inserts illustrate the characteristic asymmetric Ly α profile of C17 and ser-2.

3.1. Intrinsic Angular Diameters and Velocity Dispersion

We derive the intrinsic spatial FWHM assuming that the intrinsic, atmospheric, and instrumental FWHM add in quadrature. Based on the results of multiple measurements, we assume the 1σ observational and intrinsic FWHM measurement errors to be ~ 0.25 pix, and run 100 trials based on the observed central values. We find deconvolved FWHM for ser-1 and ser-2 of $0''.24 \pm 0''.18$ and $0''.36 \pm 0''.15$, respectively. At these redshifts, a compact galaxy with $r = 1 h_{50}^{-1}$ kpc would have an intrinsic angular diameter of $\sim 0''.27 / 0''.16$ ($\Omega = 1 / 0.1$). We note, however, that the deconvolved spatial FWHM of C17 at the Ly α line, and in the continuum redward of the line are $\sim 0''.90$ and $0''.40$, respectively. The difference is likely due to scattering of Ly α photons in the H I halo. Hence it is conceivable that the intrinsic sizes of the stellar components of ser-1 and ser-2 are $\sim 50\%$ smaller than the values obtained above, or $r \lesssim 500 h_{50}^{-1}$ pc.

We measure the instrumental resolution (for objects filling the slitlet) from the line profile of lamp spectra. We find $\text{FWHM}_{\text{instr}}^{\text{lamp}} \simeq 4.4 \text{ \AA}$. For spatially unresolved sources, we find $\text{FWHM}_{\text{instr}}^{\text{QSO}} \simeq 2.8 \text{ \AA}$, measured from the Ly α forest of the QSO SSA22 D14. The deconvolved velocity width is given by the equation,

$$\sigma_v = c \frac{\sqrt{(\text{FWHM}_{\text{obs}})^2 - (\text{FWHM}_{\text{instr}})^2}}{2.354 \lambda_{\text{obs}}}, \quad (1)$$

where c is the velocity of light. We find that σ_v is consistent with $\sim 55 \pm 8.5 \text{ km s}^{-1}$ for both serendipitous objects assuming the sources fill the slitlet (see Table 1). For unresolved sources, perhaps more appropriate for the small angular extent of these sources, $\sigma_v \lesssim 95 \text{ km s}^{-1}$. Bear in mind, however, that these values refer to the kinematics of the circum-galactic H I cloud, rather than stellar velocity dispersion, which will be less (see related discussion in §5).

3.2. Ly α Equivalent Widths

The measured continua for the serendipitously-identified Ly α -emitters were consistent with zero counts, confirming the visual impression of ser-1 and ser-2 seen in Figs. 1 and 3. Therefore, direct spectroscopic measure of the line equivalent widths is not possible. We can, however, determine lower limits on the equivalent widths using the 1σ uncertainty of the continua. The Poisson noise is the square root of the photon counts per pixel, and we denote its average per pixel by σ_{pix} . The summed noise in an aperture of width w and length l is $\sigma_{\text{pix}}\sqrt{wl}$. We have l different approximations of the value of the continuum, so the noise in the continuum σ_{cont} is,

$$\sigma_{\text{cont}} = \sigma_{\text{pix}}\sqrt{\frac{w}{l}}. \quad (2)$$

However, w is not a free parameter, but must be chosen so as to maximize the S/N . Since our sources are small, we assume their undetected continua have a Gaussian shape along a spatial cut, and that their FWHM is the same as the seeing ($\sim 0''.78$). For the Ly α line, this would imply a source of zero intrinsic extension. But since the FWHM of the continuum is probably significantly less than that of the Ly α line, as we found with SSA22 C17 (§3.1), it may in fact be a slight over-estimation of the continuum FWHM. We find the S/N ratio is maximized when $w = 1.165 \times \text{FWHM}$. For l we chose 100 pixels ($\sim 126 \text{ \AA}$). Then,

$$W_{\lambda}^{\text{obs}}(2\sigma) \gtrsim \frac{j_{\alpha}}{2\sigma_{\text{cont}}}. \quad (3)$$

For the 1σ continuum uncertainties, $\sigma_{\text{cont}} \simeq 1.7(1.4) \times 10^{-20} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ for ser-1 (ser-2). We find the 2σ lower-limits on the observed equivalent widths are $W_{\lambda}^{\text{obs}} \geq 550 \text{ \AA}$ (ser-1), and $\geq 470 \text{ \AA}$ (ser-2). The rest-frame values are tabulated in Table 1. Considering the apparent absence of N V $\lambda 1240$ and C IV $\lambda 1549$ and the low ISM velocity dispersion of these sources, the data is consistent with stellar sources, and only marginally with a possible weak AGN. We discuss this further in §4. SSA22 C17 has measured $W_{\lambda}^{\text{obs}} \simeq 35.6 \text{ \AA}$.

3.3. Star Formation Rates

Assuming the Ly α line flux is powered by star formation, the strength of this line can be used to infer a star formation rate for a young galaxy. We first estimate the H α luminosity corresponding to the measured Ly α luminosity: the low-density case B ratio Ly α /H β is approximately 25 in the absence of dust (Ferland & Osterbrock 1985), while H α /H $\beta \sim 2.8$ for hot stars ($T \simeq 1.2 \times 10^4 \text{ K}$; Osterbrock 1989). Kennicutt (1983) provides the conversion from H α luminosity to the SFR, resulting in a SFR – $\mathcal{L}_{\text{Ly}\alpha}$ relation of $\dot{M} (M_{\odot} \text{ yr}^{-1}) \simeq 10^{-42} \mathcal{L}_{\text{Ly}\alpha} (\text{ergs s}^{-1})$. This results in SFRs of $\sim 1.3/1.2 M_{\odot} \text{ yr}^{-1}$ for ser-1/ser-2, respectively. Because of the possibility of dust, or incomplete absorption of ionizing radiation by the gaseous halo, these numbers are lower limits. We believe slit losses are minimal as the lines are nearly unresolved and do not fill the slit. Line fluxes from two

other nights of observations support this conclusion. The continuum flux of the LBG SSA22 C17 at 1500 Å was used to determine the SFR of $\dot{M} = 27.7 M_{\odot} \text{ yr}^{-1}$ (Madau et al. 1998), based on a Salpeter IMF with mass range $0.1 \leq M/M_{\odot} \leq 125$. However, based on the synthetic spectra of Leitherer et al. (1995), with a continuous SFR at ~ 10 Myr of age, the SFR is ~ 1.6 times larger, or $\sim 45 M_{\odot} \text{ yr}^{-1}$. We expect that the true value is somewhere in between. The Ly α line luminosity of C17 (Table 1) would imply a SFR of $2.2 M_{\odot} \text{ yr}$.

3.4. Projected Photometric Magnitudes

We use the spectral slope of a 10^7 year old starburst with a rest equivalent width of 150 Å (Charlot & Fall 1993) to calculate the expected continua of ser-1 and ser-2. We would thus expect continuum magnitudes of $V = 27.8/28.0$ for ser-1/ser-2, or $V = 27.4/27.7$ when the Ly α line is included. The 2σ upper limits on the continua of ser-1 and ser-2 (§3.2) are only $\sim 30\%$ greater than the continua implied by $W_{\lambda}^{\text{rest}} = 150 \text{ Å}$.

4. Discussion

What is the physical origin of these isolated, high equivalent width emission lines? N V $\lambda 1240$ or C IV $\lambda 1549$ are not found in the spectra of ser-1 or ser-2. The ratio of the high-ionization line flux relative to Ly α is $\lesssim 1/15$ and $\lesssim 1/20$ for N V $\lambda 1240$ and C IV $\lambda 1549$, respectively, significantly less than ratios measured from composite AGN spectra (e.g., Stern et al. 1999). The small velocity dispersions of ser-1 and ser-2 ($\sigma_v \simeq 60 \text{ km s}^{-1}$) and the lack of associated N V $\lambda 1240$ or C IV $\lambda 1549$ emission argue against the presence of an AGN, while the high surface brightness is inconsistent with photoionization by the metagalactic flux (e.g., Bunker et al. 1998). Thus we assume the lines owe their existence to stellar sources. Below we argue that the galaxies are primeval, and attempt to fit them into a cogent picture of the evolving Universe at $z \simeq 3$.

4.1. The Production of High Equivalent Width Ly α Emission

The production of a strong Ly α line requires two things: first there must be a large column of neutral hydrogen which can intercept almost all photons with $\lambda \leq 912 \text{ Å}$, and second, Ly α radiation must escape the cloud. The large cross section of H I for Ly α photons makes Ly α extremely vulnerable to dust quenching. We consider the scenario of Tenorio-Tagle et al. (1999) in some detail as one model of the early phases of a starbursting protogalaxy. Tenorio-Tagle et al. (1999) suggest that after an initial stellar ionization of dynamically quasi-relaxed gas and attendant high equivalent width “pure” Ly α emission (i.e., a symmetric line profile), enriched stellar winds sweep up gas. The compressed gas recombines, forming a shell of neutral gas which absorbs emerging Lyman continuum radiation. Ly α radiation is then emitted and scattered within the shell.

The expansion of the compressed shell of gas results in the preferential scattering, and probable absorption, of the blue-wing of the Ly α line. The LBG SSA22 C17 (Figs. 1, 3) is a good example of this common feature of LBGs; in the insert we see a trough with center $\sim 350 \text{ km s}^{-1}$ blueward of the line center, while at $\sim 650 \text{ km s}^{-1}$ we see a second maximum. Here, perhaps, is palpable proof of a strong wind and a neutral shell, with the blueward emission line probably caused by the recombination of shocked gas associated with the expanding shell (Tenorio-Tagle et al. 1999). In this scenario, the time scale for high equivalent width Ly α line emission must be only modestly greater than the lifetime of the most massive stars (at most $\sim 10^7 \text{ yr}$).

This model is based upon a single, central starburst driving the Ly α emission — an assumption which may not generally hold in real galaxies. High-resolution VLA/optical observations of five local star-forming blue compact dwarfs reveal a history of spatially distinct star forming regions in various stages of evolution (van Zee et al. 1998). Perhaps new star-forming regions in the same galaxy could display high equivalent width Ly α emission. This effect appears to be seen in a $z = 4.04$ lensed galaxy whose star formation regions are subjected to a detailed lensing chromatography study by Bunker et al. (2000). Their color analysis shows a sequence of non-coeval star formation sites with ages ranging from zero to $\sim 100 \text{ Myr}$. “Nodes” of star formation, showing Ly α in absorption, have spectral energy distributions consistent with a burst age of $20 - 30 \text{ Myr}$. There is, however, an outlying region of very high equivalent width Ly α emission ($W_{\lambda}^{\text{rest}} > 100 \text{ \AA}$, 3σ) thought to be resonantly-scattered Ly α photons escaping from the adjacent star-forming node. Yet summed over the whole galaxy (as we would see it without the benefit of gravitational lensing), the Ly α equivalent width is modest.

We therefore suggest that, for apertures containing entire young and compact galaxies, high equivalent width stellar Ly α emission ($W_{\lambda}^{\text{rest}} \gtrsim 100 \text{ \AA}$) indicates a primeval or near-primeval galaxy. We conclude that it is very likely that ser-1 and ser-2 are primeval galaxies — by which we mean they are undergoing their first significant burst of star formation.

4.2. The Surface Density of High-Redshift Ly α -Emitters

How do ser-1 and ser-2 fit within the context of surveys of star forming galaxies at high redshift? The luminosity function (LF) of galaxies at high redshift is a useful comparative tool for our purposes. Steidel et al. (1999) find a steep faint-end slope ($\alpha \simeq -1.6$) for the observed $\mathcal{R}$ -band LF of star-forming galaxies at $\langle z \rangle \simeq 3.04$, implying that a large fraction of the UV luminosity density is produced by galaxies fainter than the spectroscopic limits of current Earth-based photometric surveys. By virtue of the association of UV luminosity with SFR, this implies that a large fraction of the star formation history is unobserved. When integrated to the apparent magnitudes expected of the continua our serendipitous emission-line galaxies ($\mathcal{R} \simeq 28 \text{ mag}$), their UV luminosity function implies number densities ~ 20 times, and integrated star formation density ~ 4 times, higher than the classical LBGs integrated to the completeness limits of their ground-based survey, $\mathcal{R} = 25 \text{ mag}$.

As we have noted, because the Ly α emission lines of such sources as our serendipitous objects are likely to be observable only for a very small fraction of 1 Gyr, the modest numbers of high equivalent width Ly α sources are tangible evidence for this many-fold larger population of slightly older, faint continuum, star forming galaxies with self-absorbed Ly α inaccessible to even the deepest spectroscopic or color selection surveys. If we are to believe that high equivalent width Ly α can only be exhibited for the first $\sim 5 \times 10^7$ yr of a galaxy’s life, and that their surface density is $\sim 3 \text{ arcmin}^{-2}$ per unit redshift, then if their distribution is uniform between $3 \leq z \leq 4$, there must be ~ 30 (50) galaxies ($\Omega = 1$ (0.1)) per square arcminute formed in that interval. This follows because there are 10 (17) intervals of 5×10^7 yr between $z = 3$ and $z = 4$. The surface density of newly formed galaxies would thus be ~ 30 times larger than the LBG surface density. One must integrate the UVLF to $\mathcal{R} = 28.5$ to achieve that surface density. Thus, available surface density estimates of primeval (high equivalent width Ly α emission) galaxies are consistent with the claim that most galaxies at $z \simeq 3$ were formed at redshifts less than $z = 4$. However, the integration of the $z \sim 4$ LF (Steidel et al. 1999) over luminosity limits identical to that at $z = 3$ shows that the implied co-moving number densities increase only slightly for Ω in the range of $0.1 \leq \Omega \leq 1.0$, apparently consistent with very low net galaxy formation. These two lines of research could be consistent if the merger rate was nearly equal to the galaxy formation rate. There is evidence that the epoch-dependent merger rate was higher in the past by $(1+z)^m$ with $m \sim 1.5-3.0$ (Windhorst et al. 1999), and that mergers were more frequent for the sub- $\mathcal{L}^*$ galaxies. However, determining the size-dependent merger rate that would be needed to leave the shape of the luminosity function unchanged while producing only a very modest increase in comoving galaxy density, is beyond the scope of this paper.

4.3. Environment

The environment of high equivalent width Ly α -emitters is of considerable interest. In their study of Lyman-break candidates in the SSA22 field, Steidel et al. (1998) display convincing evidence for redshift “spikes” at $\langle z \rangle = 3.09$, and $\langle z \rangle = 3.35$. Ser-1 lies $\sim 5000 \text{ km s}^{-1}$ from the former enhancement, a structure which they suggest is a proto-Abell cluster. The latter redshift spike, which is substantially less massive than the former, contains a quasar (SSA22 D14). Ser-2 is only $\sim 190(320)h_{50}^{-1} \text{ kpc}$ (projected), for $\Omega = 1$ (0.1), and $\lesssim 200 \text{ km s}^{-1}$ distant from the quasar, while both are on order 400 km s^{-1} beyond the apparent center of the redshift spike⁴. We note that the serendipitously-identified high equivalent width Ly α line, SSA22-S1 (Steidel et al. 1998) at $z = 3.100$, is $\sim 750 \text{ km s}^{-1}$ from the $\langle z \rangle = 3.09$ enhancement. Thus, this limited amount of data suggests that high equivalent width Ly α emitters tend to be loosely associated with density enhancements some of which may be protoclusters. Similar clustering of Ly α emitters is seen in

⁴We modeled the effect of the quasar on ser-2 by extending the observed continuum blueward of the Ly α line. The extrapolated photon flux is such that its excitation of gas around ser-2 could account for at most $\sim 3\%$ of the Ly α emission of ser-2.

studies at $z \simeq 2.4$ (Keel et al. 1999).

5. A Primeval Galaxy Model

In a speculative vein, we now consider what we have learned of primeval galaxies in the light of observations, and the modeling of associated phenomena. Our purpose is to emphasize the significance of the above results to the wider field of cosmology. Our study suggests the existence of a high volume density of tiny galaxies with high specific SFRs, which may cumulatively have rivaled that of the LBGs. We seek here to provide a galaxy formation scenario in which this is plausible, and to place this scenario in the larger context of the evolution of the universe.

In response to the very small effective radii of faint galaxies imaged by HST (e.g., Giavalisco et al. 1996b; Pascarelle et al. 1998), it has been shown that an “inside-out” galaxy formation scenario may produce galaxies with small scale-lengths (Bouwens et al. 1997; Cayón et al. 1996). A theoretical framework for this hypothesis is presented in a paper by Lin & Murray (1992), which seeks to model primeval galaxies using two alternative hypotheses of protogalactic clouds. In these models, robust self-regulating star formation occurs at a rate inversely proportional to the cooling time for hydrogen number densities greater than $n_c \sim 4 \text{ cm}^{-3}$. Their model *A* is a homogeneous cloud which collapses uniformly; the critical density is first reached at a size $R_c \simeq 6M_{11}^{1/3} \text{ kpc}$, where M_{11} is the galaxy mass in units of $10^{11} M_\odot$, creating a star-forming region whose size is inconsistent with observations of either ser-1 or ser-2. Model *B* is an isothermal cloud with a core of 500 pc, which collapses in an inside-out manner according to the Shu (1977) self-similar collapse model, and reaches the density n_c at $R_c \simeq 1M_{11}^{1/2} R_{100}^{-1/2} \text{ kpc}$, where R_{100} is the outer radius of the cloud in units of 100 kpc. Model B predicts sizes that are very close to the constrained sizes of our serendipitous emission line regions, and makes their comparison with the model appear quite promising. It also predicts that protogalaxies will have a relatively constant bolometric luminosity for a period of $\sim 1.7 \text{ Gyr}$. In their published simulation, the bolometric luminosity $\log \mathcal{L} \sim 43.5 (\text{ergs}^{-1})$, though the scale-free nature of isothermal spheres could be expected to allow the somewhat more modest luminosities observed here. We shall see that this luminosity is consistent with a SFR of $\sim 5.7 M_\odot \text{ yr}^{-1}$.

An isothermal cloud which is bounded by an external medium of pressure P_{ext} may achieve a hydrostatic equilibrium if the ratio of the external to the internal pressure is less than 14.3 (Shu 1977). This cloud, known as a Bonnor-Ebert sphere, has a just-critical mass $M_{\text{crit}} \simeq 0.8a^4 G^{-3/2} P_{\text{ext}}^{-1/2}$, where a is the sound speed. This mass may be consistent with galactic masses when the intergalactic pressure is modest (i.e., $P_{\text{ext}} \lesssim 1 \times 10^{-19} \text{ dyne cm}^{-2}$). Such a sphere would become super-critical, and collapse from the inside out if it were part of a general over-density which had pulled away from the Hubble flow and begun to collapse. Linear theory shows that the turn-around radius moves outward at a rate of $\sim$ half the Hubble flow (Davis et al. 1980), so that we could expect the location of high equivalent width Ly α emission to be somewhat within the turn-around radius, and to move outward with time. A more extensive discussion of this subject will be reserved for a

future publication.

In order to link the model with observations, we made a rough simulation of a galaxy undergoing a constant SFR over a period of up to 1 Gyr. We used a Salpeter IMF ($\alpha = -2.35$) with lower mass limit of $0.1 M_{\odot}$, and main sequence lifetimes and luminosities (Tables 3-6 and 3-9) from Mihalas & Binney (1981), modeling them as black bodies, which is sufficiently accurate for metal-free, non-evolved stars. Unless otherwise stated, our upper mass limit is $125 M_{\odot}$, and we assume that the SFR is $1.2 M_{\odot} \text{ yr}^{-1}$. The Ly α photon emission rate of a primeval galaxy is taken to be 2/3 of the stellar emission of photons more energetic than 13.6 eV, according to the case B recombination (Spitzer 1978). We assume that 80% of all newly introduced gas (i.e., over and above the $\sim 10^7 M_{\odot}$ of gas of density n_c within the central 500 pc) is transformed into stars immediately.

The continuum measured in the 75 Å region around the Ly α emission line, together with the Ly α luminosity, is used to calculate the equivalent width. We tested the derived equivalent widths against the dust-free calculations of Charlot & Fall (1993) for continuous star formation and upper mass limits of 80 and 120 $M_{\odot}$, and found values that were quite consistent with values attained by interpolating between the bracketing Salpeter slopes which they employed. We find that for a 10 Myr old galaxy, $W_{\lambda}^{\text{rest}} = 160/200/250 \text{ Å}$ respectively for 80/100/120 $M_{\odot}$ upper limits. These derived values drop to $\sim 125/155/200 \text{ Å}$ by 100 Myr. The relation of the resulting Ly α line luminosity to the SFR is in agreement with the case B predictions. The model predicts a continuum in the observed visual band of $\sim 1.3 \times 10^{-20} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Å}^{-1}$ for a $z = 3$ galaxy with $\dot{M} = 1.2 M_{\odot} \text{ yr}^{-1}$. This is within the 1σ non-detection limits of the continua of ser-1 and ser-2 ($1.7/1.4 \times 10^{-20} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Å}^{-1}$).

For an upper mass limit of 100 $M_{\odot}$, and $\dot{M} = 1.0 M_{\odot} \text{ yr}^{-1}$, the bolometric luminosity at 10 Myr is $\sim 5.5 \times 10^{42} \text{ ergs s}^{-1}$. Thus, the luminosity of the Lin & Murray (1992) model B corresponds to $\dot{M} \simeq 5.7 M_{\odot} \text{ yr}^{-1}$. We calculate the mass to bolometric luminosity ratio of our model by adding the mass in stars and dissipated gas, dividing it by the luminosity. The dark matter does not participate in the cloud's collapse since it is not pressure supported. It will be compressed somewhat by the gradually increasing concentration of baryons in the center, however, on the whole, the DM halo should have little effect on the velocity dispersion of stars formed out of dissipated gas in the first few tens of Myr since they are formed well within the core of the isothermal sphere. We find that the calculated $\mathcal{M}/\mathcal{L} \simeq 0.01, 0.05, \text{ and } 0.37$ for $\log t = 7, 8, \text{ and } 9$ (Gyr) respectively. We note that compact narrow emission line galaxies, which appear to be likely lower-redshift counterparts, have mass-to-luminosity ratios, $\mathcal{M}/\mathcal{L} \approx 0.1$ (Phillips et al. 1997).

The expected stellar velocity dispersion, $\sigma_s = \sqrt{GM_{\text{stars+gas}}/r}$, implies dispersions of model galaxies of from $\sim 14 / 37 /$, and 110 km s^{-1} for $\log t = 7, 8, \text{ and } 9$ (yr), assuming a radius of 500 pc. The low σ_s for the 10 Myr old galaxy is actually reasonable, in view of the finding that the very large cluster in the blue compact dwarf NGC 1705 (Ho & Filippenko 1996) has $\sigma_s \sim 11 \text{ km s}^{-1}$. A low σ_s is also not in conflict with our deconvolved velocity dispersions of $\sigma_v \sim 60 \text{ km s}^{-1}$ (§3.1) since the latter is a measure of both the turbulence of gas (from infall and stellar winds) which give

rise to the Ly α resonance line, as well as the possibly spatially resolved structure within the slitlet. On the basis of the observed σ_v and deconvolved radius r of the LBG C17 (Table 1), we find that the galaxy mass is $M_{\text{LBG}} = 4.3 \times 10^9 M_{\odot}$. This galaxy would take ~ 1.2 Gyr to form at the SFR derived for ser-1 or ser-2. However, between $z = 3.5$ and $z = 3.0$, there are only 0.265/0.473 Gyr for $\Omega = 1.0/0.1$. Thus, LBGs would not be formed in a reasonable amount of time from low-SFR objects if SFRs remain constant. Below, we consider the possibility that the SFR may increase with time

Our simple galaxy model has assumed that $\dot{M}$ is constant. However, a realistic isothermal sphere will have a baryonic core, assumed here to be ~ 500 pc. In this case the infall rate will *not* be constant in time, but given a sound speed of 10 km s^{-1} , may increase during the first $\sim 5 \times 10^7$ yr. In addition, the negligible metallicity expected of primordial gas will make cooling times, and the Jeans mass, larger⁵ during the first few tens of Myr of a galaxy’s life. Thus the effect of metals ejected from evolved stars in aiding cooling, and of dust in catalyzing the production of molecules, is to compound the effect of increasing infall; the SFR and its efficiency can be expected to rise substantially by a few tens of Myr.

On the other hand, recall that the high equivalent width of the Ly α emission line of primeval galaxies is expected to be quenched over essentially the same time scale. This means that the nature, and fate of these pre-LBG/post-primeval galaxies will be very difficult to learn. With an enhanced SFR, the post-primeval galaxies could plausibly grow by a factor of 10-40 into LBGs in a few tenths of a Gyr or, by virtue of competition from neighbors, have their growth truncated while still in the dwarf stage, or even be cannibalized; environment may have a dominant role to play.

While the mass and SFRs of our purportedly primeval galaxies are much smaller than that of LBGs, the *specific* SFR of the former is on average 10 or 20 times greater than the latter. The large numbers of small, young galaxies which are thought to have been formed (§4.2) in the interval $3 \lesssim z \lesssim 4$ might thus be expected to provide a substantial fraction of the star formation rate density. In fact, the steep faint end slope of the $z \sim 3$ UV luminosity function of Steidel et al. (1999) implies that up to 40% of the SFR density may be supplied by galaxies fainter than $\mathcal{R} = 27$ ($\mathcal{L} \sim 0.1$) but brighter than the projected continuum magnitudes of our serendipitous sources.

In addition to the formation of a galactic bulge according to the Lin-Murray process, it is expected to be accompanied by the formation of the halo, as infalling clumps of gas dissipatively interact with the already accreted gas (Binney 1976; Manning 1999).

⁵It has been suggested (Lin & Murray 1992) that the IMF of primeval galaxies may be heavily weighted in high-mass stars, which could result in rest Ly α equivalent widths significantly larger than $150 \text{ \AA}$. The truth of this contention could be tested if spectroscopic observations deep enough to detect the continua were made.

6. Conclusions

We draw the following conclusions from this study:

- We suggest that the combination of high equivalent widths, low velocity dispersions, and small intrinsic sizes seen in the Ly α lines of these two galaxies can be well-explained as their being extremely low-metallicity galaxies undergoing an initial burst of star formation—hence, primeval galaxies. These characteristics, markedly different from earlier expectations, suggests an inside-out, rather than a monolithic collapse formation mechanism.
- Our data, which imply that most galaxies at $z = 3$ may have been formed at $z \leq 4$, stand in apparent conflict with the $z = 3$ and 4 UVLFs (Steidel et al. 1999), which suggest that the comoving galactic density remained relatively constant during this interval. These may be reconciled, however, with a large merger rate that preserves the faint-end slope of the luminosity function.
- The agreement of the angular size, and the projected total luminosity of these emission line galaxies, with the predictions of the inside-out collapse of a nearly-isothermal H I halo (Lin & Murray 1992) suggests that the latter is a promising basis for modeling galaxy formation. Our discussion has suggested that a rising star formation rate with a time-scale of $\sim 5 \times 10^7$ yr is plausible for many post-primeval galaxies.
- The isothermal collapse model requires that protogalactic clouds form in relatively isolated regions. However, their collapse may be stimulated by the gradually increasing pressures of a protocluster environment, thus accounting for the apparent weak clustering on large scales noted of high-, as well as low-redshift Ly α emission line galaxies.

We thank C. Leitherer, and C. C. Steidel, for useful discussions. We are grateful to our referee, Dr. R. A. Windhorst, for valuable suggestions. We also thank A. Dey for assistance with the observations. We are grateful for the support of NSF grant AST 95-28536.

References

- Binney, J. 1976, MNRAS, 177, 19
- Bouwens, R. J., Cayón, L., & Silk, J. 1997, ApJ, 489, L21
- Bunker, A. J., Marleau, F. R., & Graham, J. R. 1998, AJ, 116, 2086
- Bunker, A. J., Moustakas, L. A., & Davis, M. 2000, ApJ, in press
- Cayón, L., Silk, J., & Charlot, S. 1996, ApJ, 467, L53

- Charlot, S. & Fall, S. M. 1993, ApJ, 451, 580
- Cowie, L. & Hu, E. M. 1998, AJ, 115, 1319
- Davis, M., Tonry, J., Huchra, J., & Latham, D. W. 1980, ApJ, 238, L113
- Dey, A., Spinrad, H., Stern, D., Graham, J. R., & Chaffee, F. 1998, ApJ, 498, L93
- Ferland, G. J. & Osterbrock, D. E. 1985, ApJ, 289, 105
- Giavalisco, M., Koratkar, A., & Calzetti, D. 1996a, ApJ, 466, 831
- Giavalisco, M., Steidel, C. C., & Macchetto, D. 1996b, ApJ, 470, 189
- Ho, L. C. & Filippenko, A. V. 1996, ApJ, 472, 600
- Hu, E. M., Cowie, L. L., & McMahon, R. G. 1998, ApJ, 502, 99
- Hu, E. M., McMahon, R. G., & Cowie, L. L. 1999, ApJ, 522, 9
- Keel, W. C., Cohen, S. H., Windhorst, R. A., & Waddington, I. 1999, AJ, 118, 2547
- Kennicutt, R. 1983, ApJ, 282, 54
- . 1992, ApJ, 388, 310
- Kunth, D., Mas-Hesse, J. M., Terlevich, E., Terlevich, R., & Fall, S. M. 1998a, A&A, 334, 11
- Kunth, D., Terlevich, E., Terlevich, R., & Tenorio-Tagle, G. 1998b, in *Dwarf Galaxies and Cosmology*, ed. T. X. Thuan (Gif-sur-Yvette: Editions Frontieres), in press
- Legrand, F., Kunth, D. F., Mas-Hesse, J. M., & Lequeus, J. 1997, A&A, 326, 929
- Leitherer, C., Carmelle, R., & Heckman, T. M. 1995, ApJS, 99, 173
- Lin, D. N. C. & Murray, S. D. 1992, ApJ, 394, 523
- Lowenthal, J. D. et al. 1997, ApJ, 481, 673
- Madau, P., Pozzetti, L., & Dickinson, M. E. 1998, ApJ, 498, 106
- Manning, C. 1999, ApJ, 518, 226
- Massey, P. & Gronwall, C. 1990, ApJ, 358, 344
- Mihalas, D. & Binney, J. 1981, *Galactic Astronomy* (San Francisco: W. H. Freeman and Company)
- Oke, J. B. et al. 1995, PASP, 107, 375
- Osterbrock, D. 1989, *The Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Mill Valley: Univ. Science Books)

- Pascarelle, S. M., Windhorst, R. A., & Keel, W. C. 1998, *AJ*, 116, 2659
- Pascarelle, S. M., Windhorst, R. A., Keel, W. C., & Odewahn, S. C. 1996, *Nature*, 383, 45
- Phillips, A. C., Guzman, R., Gallego, J., Koo, D. C., Lowenthal, J. D., Vogt, N. P., Faber, S. M., & Illingworth, G. D. 1997, *ApJ*, 489, 543
- Pritchett, C. J. 1994, *PASP*, 106, 1052
- Shu, F. H. 1977, *ApJ*, 214, 488
- Spitzer, L. 1978, *Physical Processes in the Interstellar Medium* (New York: Wiley)
- Steidel, C. S., Adelberger, K. L., Dickinson, M., Giavalisco, M., Pettini, M., & Kellogg, M. 1998, *ApJ*, 492, 428
- Steidel, C. S., Adelberger, K. L., Giavalisco, M., Dickinson, M., & Pettini, M. 1999, *ApJ*, 519, 1
- Steidel, C. S., Giavalisco, M., Dickinson, M., & Adelberger, K. L. 1996, *AJ*, 112, 352
- Steidel, C. S. & Hamilton, D. 1992, *AJ*, 104, 941
- Stern, D., Bunker, A. J., Spinrad, H., & Dey, A. 2000, *ApJ*, in press
- Stern, D., Dey, A., Spinrad, H., Maxfield, L. M., Dickinson, M. E., Schlegel, D., & González, R. A. 1999, *AJ*, 117, 1122
- Stockton, A. & Ridgway, S. E. 1998, *AJ*, 115, 1340
- Tenorio-Tagle, G., Silich, S. A., Kunth, D., Terlevich, E., & Terlevich, R. 1999, *MNRAS*, 309, 332
- Terlevich, E., Díaz, A. I., Terlevich, R., & García, M. L. 1993, *MNRAS*, 260, 3
- Thompson, D. & Djorgovski, S. G. 1995, *AJ*, 110, 982
- van Zee, L., Skillman, E. D., & Salzer, J. J. 1998, *AJ*, 110, 963
- Weymann, R., Stern, D., Bunker, A. J., Spinrad, H., Chaffee, F., Thompson, R., & Storrie-Lombardi, L. 1998, *ApJ*, 505, L95
- Windhorst, R. A., Cohen, S. H., & Waddington, I. 1999, in *After the Dark Ages: When Galaxies were Young*, ed. S. S. Holt & E. P. Smith, Vol. 470 (New York: AIP), 202